

Introduction to Statistics

lab 8

Johan A. Elkink
jos.elkink@ucd.ie

5 November 2018

Data

For this lab we will continue to make use of the Irish National Election Study (INES) data from the 2007 general election. See Lab 6 for download and subset instructions.

t-tests

As will be clear in this and the next lecture, there are three types of *t*-tests, which can be executed as follows:

- One-sample *t*-test, where you compare the mean of one variable against a fixed value. For example, to test whether the mean of a variable **v0190** differs from 5:
SPSS: T-TEST TESTVAL = 5 /VARIABLES = v0190.
R: `t.test(v0190, mu = 5)`
Stata: `ttest v0190 == 5`
- Paired-samples *t*-test, where you compare the mean on two variables for the same individuals (e.g. a test score before and after a class). For example, to test whether the means of two variables, **v0190** and **v0191**, for the same individuals differ:
SPSS: T-TEST PAIRS = v0190 v0191.
R: `t.test(v0190, v0191, paired = TRUE)`
Stata: `ttest v0190 == v0191`
- Independent-samples *t*-test, where you compare the mean on the same variable for two different groups. For example, **v0190** depending on whether a variable **union** is 1 or 0:
SPSS: T-TEST GROUPS = union(0,1) /VARIABLES = v0190.
R: `t.test(v0190 ~ union)`
Stata: `ttest v0190, by(union)`

For the following statements, formulate the null hypothesis and the alternative hypothesis, perform the appropriate *t*-test, and formulate the conclusion from the test:

1. Voters are more likely to vote Labour (**v0190**) than they are to vote Sinn Fein (**v0192**).
2. Union members are more likely to vote Labour (**v0190**) than are non-union members (**v0936**).
3. Voters in Ireland tend to the political right (i.e. **v0239** is greater than 5 on average).
4. Fianna Fáil voters are more right-wing (**v0239**) than Fine Gael voters (**v0195**), on average.
5. Voters like Enda Kenny (**v0522**) more than they like Pat Rabbitte (**v0524**).¹
6. *Extra challenge*: Voters who agree that there should be very strict limits on immigration (from “slightly agree” to “strongly agree”) (**v0247**) are more likely to vote for Sinn Fein (**v0192**) than the others.

Attitudes towards abortion among younger voters

1. Construct a new variable **age** which is calculated on the basis of **v0906**. Remember that the year of the survey is 2007. SPSS users will need the COMPUTE command – cf. how you calculated the logarithmically transformed variables (see Lab 2). Stata users will use gen.
2. Produce a frequency table of attitude towards abortion (**v0266**) to check whether missing values and numerical values are correctly coded.²
3. Produce a scatter plot of attitude towards abortion by **age**. Add a regression line.
4. Regress attitude towards abortion on age.
5. Recode age into a variable **young**, which is 1 if **age** is less than 30 and 0 otherwise.
6. Regress attitude towards abortion on the young dummy variable. (Note comments in Lab 6.)
7. Perform a *t*-test whether the mean on abortion is different for young versus old. Compare the result to the regression result.

¹R users might have to check whether it is properly coded as a numerical variable.

²R users need to include the option `exclude = NULL` in the `table()` command to include the missing values.